

Autosomal dominant nocturnal frontal lobe epilepsy

Description

Autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE) is an uncommon form of epilepsy that runs in families. This disorder causes seizures that usually occur at night (nocturnally) while an affected person is sleeping. Some people with ADNFLE also have seizures during the day.

The seizures characteristic of ADNFLE tend to occur in clusters, with each one lasting from a few seconds to a few minutes. Some people have mild seizures that simply cause them to wake up from sleep. Others have more severe episodes that can include sudden, repetitive movements such as flinging or throwing motions of the arms and bicycling movements of the legs. The person may get out of bed and wander around, which can be mistaken for sleepwalking. The person may also cry out or make moaning, gasping, or grunting sounds. These episodes are sometimes misdiagnosed as nightmares, night terrors, or panic attacks.

In some types of epilepsy, including ADNFLE, a pattern of neurological symptoms called an aura often precedes a seizure. The most common symptoms associated with an aura in people with ADNFLE are tingling, shivering, a sense of fear, dizziness (vertigo), and a feeling of falling or being pushed. Some affected people have also reported a feeling of breathlessness, overly fast breathing (hyperventilation), or choking. It is unclear what brings on seizures in people with ADNFLE. Episodes may be triggered by stress or fatigue, but in most cases the seizures do not have any recognized triggers.

The seizures associated with ADNFLE can begin anytime from infancy to mid-adulthood, but most begin in childhood. The episodes tend to become milder and less frequent with age. In most affected people, the seizures can be effectively controlled with medication.

Most people with ADNFLE are intellectually normal, and there are no problems with their brain function between seizures. However, some people with ADNFLE have experienced psychiatric disorders (such as schizophrenia), behavioral problems, or intellectual disability. It is unclear whether these additional features are directly related to epilepsy in these individuals.

Frequency

ADNFLE appears to be an uncommon form of epilepsy; its prevalence is unknown. This

condition has been reported in more than 100 families worldwide.

Causes

Mutations in the *CHRNA2*, *CHRNA4*, and *CHRNA2* genes can cause ADFLE. These genes provide instructions for making different parts (subunits) of a larger molecule called a neuronal nicotinic acetylcholine receptor (nAChR). This receptor plays an important role in chemical signaling between nerve cells (neurons) in the brain.

Communication between neurons depends on chemicals called neurotransmitters, which are released from one neuron and taken up by neighboring neurons. Researchers believe that mutations in the *CHRNA2*, *CHRNA4*, and *CHRNA2* genes affect the normal release and uptake of certain neurotransmitters in the brain. The resulting changes in signaling between neurons likely trigger the abnormal brain activity associated with seizures.

The seizures associated with ADFLE begin in areas of the brain called the frontal lobes. These regions of the brain are involved in many critical functions, including reasoning, planning, judgment, and problem-solving. It is unclear why mutations in the *CHRNA2*, *CHRNA4*, and *CHRNA2* genes cause seizures in the frontal lobes rather than elsewhere in the brain. Researchers are also working to determine why these seizures occur most often during sleep.

The genetic cause of ADFLE has been identified in only a small percentage of affected families. In some cases, a gene other than those that make up the nAChR are involved. In the remaining families, the cause of the condition is unknown. Researchers are searching for other genetic changes, including mutations in other subunits of nAChR, that may underlie the condition.

[Learn more about the genes associated with Autosomal dominant nocturnal frontal lobe epilepsy](#)

- *CHRNA2*
- *CHRNA4*
- *CHRNA2*
- *KCNT1*

Inheritance

This condition is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to raise the risk of developing epilepsy. About 70 percent of people who inherit a mutation in the *CHRNA2*, *CHRNA4*, or *CHRNA2* gene will develop seizures. In most cases, an affected person has one affected parent and other relatives with the condition. Other cases are described as sporadic, which means an affected person has no family history of the disorder.

Other Names for This Condition

- ADNFLE

Additional Information & Resources

Genetic Testing Information

- Genetic Testing Registry: Epilepsy, nocturnal frontal lobe, type 1 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1838049/>)
- Genetic Testing Registry: Epilepsy, nocturnal frontal lobe, type 2 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1864125/>)
- Genetic Testing Registry: Epilepsy, nocturnal frontal lobe, type 3 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1854335/>)
- Genetic Testing Registry: Epilepsy, nocturnal frontal lobe, type 4 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1835905/>)

Genetic and Rare Diseases Information Center

- Autosomal dominant nocturnal frontal lobe epilepsy (<https://rarediseases.info.nih.gov/diseases/11918/autosomal-dominant-nocturnal-frontal-lobe-epilepsy>)

Patient Support and Advocacy Resources

- Disease InfoSearch (<https://www.diseaseinfosearch.org/>)
- National Organization for Rare Disorders (NORD) (<https://rarediseases.org/>)

Research Studies from ClinicalTrials.gov

- ClinicalTrials.gov (<https://clinicaltrials.gov/ct2/results?cond=%22Epilepsy%2C+Frontal+Lobe%22+OR+%22autosomal+dominant+nocturnal+frontal+lobe+epilepsy%22>)

Catalog of Genes and Diseases from OMIM

- EPILEPSY, NOCTURNAL FRONTAL LOBE, 1 (<https://omim.org/entry/600513>)
- EPILEPSY, NOCTURNAL FRONTAL LOBE, 2 (<https://omim.org/entry/603204>)
- EPILEPSY, NOCTURNAL FRONTAL LOBE, 3 (<https://omim.org/entry/605375>)
- EPILEPSY, NOCTURNAL FRONTAL LOBE, 4 (<https://omim.org/entry/610353>)

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28autosomal+dominant+nocturnal+frontal+lobe+epilepsy%5BTIAB%5D%29+OR+%28adnfl%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>)

References

- Combi R, Dalprà L, Tenchini ML, Ferini-Strambi L. Autosomal dominant nocturnal frontal lobe epilepsy--a critical overview. *J Neurol*. 2004 Aug;251(8):923-34. Review. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15316796>)
- De Marco EV, Gambardella A, Annesi F, Labate A, Carrideo S, Forabosco P, Civitelli D, Candiano IC, Tarantino P, Annesi G, Quattrone A. Further evidence of genetic heterogeneity in families with autosomal dominant nocturnal frontal lobe epilepsy. *Epilepsy Res*. 2007 Apr;74(1):70-3. Epub 2007 Feb 26. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17324557>)
- di Corcia G, Blasetti A, De Simone M, Verrotti A, Chiarelli F. Recent advances on autosomal dominant nocturnal frontal lobe epilepsy: "understanding the nicotinic acetylcholine receptor (nAChR)". *Eur J Paediatr Neurol*. 2005;9(2):59-66. Review. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15843070>)
- Hayman M, Scheffer IE, Chinvarun Y, Berlangieri SU, Berkovic SF. Autosomal dominant nocturnal frontal lobe epilepsy: demonstration of focal frontal onset and intrafamilial variation. *Neurology*. 1997 Oct;49(4):969-75. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9339675>)
- Kurahashi H, Hirose S. Autosomal Dominant Nocturnal Frontal Lobe Epilepsy. 2002 May 16 [updated 2018 Mar 15]. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Mirzaa GM, Amemiya A, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2022. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1169/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301348>)
- Marini C, Guerrini R. The role of the nicotinic acetylcholine receptors in sleep-related epilepsy. *Biochem Pharmacol*. 2007 Oct 15;74(8):1308-14. Epub 2007 Jun 23. Review. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17662253>)
- Oldani A, Zucconi M, Asselta R, Modugno M, Bonati MT, Dalprà L, Malcovati M, Tenchini ML, Smirne S, Ferini-Strambi L. Autosomal dominant nocturnal frontal lobe epilepsy. A video-polysomnographic and genetic appraisal of 40 patients and delineation of the epileptic syndrome. *Brain*. 1998 Feb;121 (Pt 2):205-23. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9549500>)
- Provini F, Plazzi G, Tinuper P, Vandi S, Lugaresi E, Montagna P. Nocturnal frontal lobe epilepsy. A clinical and polygraphic overview of 100 consecutive cases. *Brain*. 1999 Jun;122 (Pt 6):1017-31. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10356056>)
- Scheffer IE, Bhatia KP, Lopes-Cendes I, Fish DR, Marsden CD, Andermann E, Andermann F, Desbiens R, Keene D, Cendes F, et al. Autosomal dominant nocturnal frontal lobe epilepsy. A distinctive clinical disorder. *Brain*. 1995 Feb;118 (

Pt 1):61-73. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/7895015>)

Page last updated on 18 August 2020

Page last reviewed: 1 April 2009